

РОЛЬ ГЕОМЕТРИИ В СОВРЕМЕННОЙ ФИЗИКЕ

С.Г Рубин

В работе обсуждается роль дополнительных измерений. Демонстрируется, что идея дополнительного пространства в комбинации с теорией гравитации с высшими производными является подходящим инструментом для решения фундаментальных проблем современной физики.

Ключевые слова: Дополнительное пространство, размерность, гравитация, тонкая настройка.

UDC 514.822

ON CONFORMAL KILLING AND HARMONIC FORMS ON RIEMANNIAN SYMMETRIC SPACES

S.E. Stepanov¹, T.V. Dmitrieva²

¹ s.e.stepanov@mail.ru; Financial University at the Government of the Russian Federation

² ztv77@mail.ru; Russian State Social University

We prove that there are no non-zero closed and coclosed conformal Killing L^2 -forms on a complete, simply connected and irreducible symmetric space with negative scalar curvature. We also prove that there are no non-zero harmonic forms on a complete, simply connected and irreducible symmetric space with positive scalar curvature.

Keywords: Riemannian symmetric space, harmonic forms, closed and coclosed conformal Killing forms.

Conformal Killing forms have been defined on Riemannian manifolds more than forty-five years ago by Tachibana (see [1]) as a natural generalization of conformal Killing vector fields. Surveys of the publications on these forms can be found in the introduction to our last paper [2].

A Riemannian globally symmetric space of *non-compact type* (M, g) is complete and also (M, g) has a nonpositive sectional curvature. We also know that a Riemannian symmetric space has nonpositive (resp. non-negative) curvature operator if and only if it has nonpositive (resp. non-negative) sectional curvature (see [3]). Note that symmetric spaces of non-compact type are non-compact. After the above remarks, the assertion of the following theorem becomes obvious.

Theorem 1. *Let (M, g) be an n -dimensional ($n \geq 3$) complete, simply connected symmetric space. If (M, g) is irreducible and its scalar curvature is negative, then there are no non-zero closed and coclosed conformal Killing L^2 -forms of degree p ($1 \leq p \leq n-1$) on (M, g) .*

Theorem 2. *Let (M, g) be an $2p$ -dimensional ($p \geq 1$) complete, simply connected symmetric space. If (M, g) is irreducible and its scalar curvature is negative, then there are no non-zero conformal Killing L^2 -forms of degree p on (M, g) .*

It is well known that a Riemannian globally symmetric space of *compact type* (M, g) is compact and also (M, g) has a nonpositive sectional curvature (see [3]). Then the following theorem holds.

Theorem 3. *Let (M, g) be an n -dimensional ($n \geq 2$) complete, simply connected symmetric space. If (M, g) is irreducible and its scalar curvature is positive, then there are no non-zero harmonic p -forms ($1 \leq p \leq n - 1$) on (M, g) .*

Our work was supported by RBRF grant 16-01-00756-a (Russia).

References

1. Tachibana S. *On conformal Killing tensor in a Riemannian space* // Tohoku Math. Journal – 1969. – V. 21. – С. 56–64.
2. Stepanov S. E., Tsyganok I. I., Dmitrieva T. V. *Harmonic and conformal Killing forms on complete Riemannian manifold* // Russian Math. – 2017. – V. 61. – № 3. – С. 44–48.
3. Duchesne B. *Infinite dimensional Riemannian symmetric spaces with fixed-sing curvature operator* // Ann. Inst. Fourier, Grenoble – 2015. – V. 65. – № 1. – С. 211–244.

О КОНФОРМНО КИЛЛИНГОВЫХ И ГАРМОНИЧЕСКИХ ФОРМАХ НА РИМАНОВЫХ СИММЕТРИЧЕСКИХ ПРОСТРАНСТВАХ

С.Е. Степанов, Т.В. Дмитриева

Мы доказываем, что не существует ненулевых замкнутых и козамкнутых конформно киллинговских L^2 -форм на полном односвязном неприводимом римановом симметрическом многообразии с отрицательной скалярной кривизной. Также доказываем, что не существует гармонических L^2 -форм на полном односвязном неприводимом римановом симметрическом многообразии с положительной скалярной кривизной.

Ключевые слова: Риманово симметрическое пространство, гармонические формы, замкнутые и козамкнутые конформно киллинговые формы.

UDC 514.822

ON INFINITESIMAL HARMONIC TRANSFORMATIONS ON COMPLETE RIEMANNIAN MANIFOLDS

S.E. Stepanov¹, I.I. Tsyganok²

¹ s.e.stepanov@mail.ru; Financial University at the Government of the Russian Federation

² i.i.tsyganok@mail.ru; Financial University at the Government of the Russian Federation

We prove that every infinitesimal harmonic transformation with finite kinetic energy on a complete Riemannian manifold with non-positive Ricci curvature is parallel. We apply this fact to the Ricci soliton theory.

Keywords: Complete Riemannian manifold, infinitesimal harmonic transformation, Ricci soliton.

A vector field ξ on a complete Riemannian manifold (M, g) is called an infinitesimal harmonic transformation if ξ generates a flow which is a local one-parameter group of harmonic transformations [1].

The kinetic energy of the flow (see [2, p. 2]) on a complete Riemannian manifold (M, g) generated by a complete vector field ξ is determined by the following equation $E(\xi) = \frac{1}{2} \int_M \|\xi\|^2 dVol_g$. Using the above definition we can prove the following theorem.